

1 **Supplementary Material for**
2 **Atmospheric and Oceanic Origins of Tropical Precipitation Variability**

3

4 Jie He¹, Clara Deser² & Brian J. Soden³

5

6

7 1. Princeton University, and NOAA/Geophysical Fluid Dynamics Laboratory, Princeton,

8

New Jersey

9

2. National Center for Atmospheric Research, Boulder, Colorado

10

3. Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami,

11

Florida

12

13

14

15

16

17

18 *Corresponding author's address:*

19

Jie He

20

Geophysical Fluid Dynamics Laboratory

21


Princeton University

22

201 Forrestal Road, Princeton, NJ, 08540, USA

23

E-mail: Jie.He@noaa.gov



24

25 Figure S1. Differences in annual mean surface temperature climatology between (a)

26 Coupled and ClimSST, (b) Coupled and FullSST and (c) Coupled and SlabOcean.

27 Contours show the climatological surface temperature climatology with an interval of 3K.

28 Differences in DJF and JJA climatologies show similar patterns. Areas where the

29 difference in variance is not significant at the 99% level based on the student t-test are

30 stippled.


31

32

33

34

35




36

37 Figure S2. Mean (contours) and standard deviation (color shading) of daily precipitation
 38 anomalies from (a) the Coupled run, (b) the FullSST run and (c) their difference. Results
 39 from the Coupled run are calculated as the average of 18 non-overlapping 100-year
 40 segments, whereas results for the FullSST run are calculated using the available 100
 41 years. Contour interval is 3 mm/day starting at 3 mm/day in (a) and (b), and 0.3 mm/day
 42 in (c) with dashed lines indicating negative values. Stippling in (c) indicates that the
 43 difference in variance does not pass the 99% significance level based on the f-test. Note
 44 that the FullSST run is forced with monthly mean SST anomalies. The small difference in
 45 variance between the two runs (which matches the difference in climatology) indicates

46 that the lack of two-way coupling and the lack of sub-monthly SST variability do not
 47 substantially affect the simulation of precipitation variance.

48

49



50

51 Figure S3. Regional average surface temperature power spectra from the Coupled run.

52 The location of the regions is shown as the purple boxes in Figure 2g in the main text.


53 The spectra are calculated as the average of partially overlapping 100-year segments from

54 the daily output.

55

56

57




58

59 Figure S4. Fractional difference in precipitation standard deviation (color shading)
 60 between (a) Coupled and ClimSST, (b) SlabOcean and ClimSST and (c) Coupled and
 61 SlabOcean, using (left) 10-year high pass, (middle) 10 to 50-year band pass and (right)
 62 50-year low pass yearly DJF precipitation anomalies. The fractional difference is shown
 63 as a percentage relative to the Coupled standard deviation. DJF mean precipitation
 64 climatology from the Coupled run is plotted as contours. Contour interval is 3 mm/day
 65 starting at 3 mm/day.

66


67



68

69 Figure S5. The same as Figure S4, except for JJA.

70



71

72 Figure S6. Pointwise correlation between DJF anomalies of precipitation and surface
 73 temperature in (a) the Coupled run and (b) observations and between DJF anomalies of
 74 precipitation and the negative of the Laplacian of surface temperature in (c) the Coupled
 75 run and (d) observations. The observed surface temperature is taken from the merged
 76 Hadley-NOAA/Optimal Interpolation SST (Hurrell et al. 2008) instead of GISS because
 77 the latter only provides surface temperature anomalies, which alone does not yield the
 78 Laplacian of surface temperature without information about the mean climatology. The

79 observations span 1979/02 to 2012/01. The Laplacian is calculated using spherical
80 harmonics, and all observations are interpolated to the CAM model grid before
81 calculating the Laplacian. The DJF mean precipitation climatology is plotted as contours.
82 Contour interval is 3 mm/day starting at 3 mm/day. Stippling indicates that the linear
83 correlation between precipitation and surface temperature or between precipitation and
84 the Laplacian of surface temperature is not significant at the 99% level based on the two-
85 sided t-test. A comparison between (a, b) and (c, d) indicates that monthly precipitation
86 anomalies are more closely related to local SST rather than local SST gradient.

87

88


89

90

91


92

93



94

95 Figure S7. The same as Fig. S6, except for JJA.




96

97 Figure S8. Precipitation (shading) and surface temperature (contour) correlation with
98 Nino 3.4 SST anomaly using yearly DJF (top) and JJA (bottom) output from the Coupled
99 run. The Nino 3.4 region is defined as the area from 5S-5N and 170-120W. Contour
100 interval is 0.2. Dashed contours represent negative values. The zero contour is thickened.

101

102

103



104

105 Figure S9. The same as Figure 12 in the main text, except for EOF2 instead of EOF1.

106

107


108

109

110

111


112



113

114 Figure S10. The same as Figure 14 in the main text, except for North Pacific (outlined by

115 the red box) instead of South Pacific.



116

117 Figure S11. Linear correlation of surface temperature (shading) and SLP (contour)

118 anomalies with the leading PC of surface temperature anomalies in the tropical Pacific

119 region, which is marked by the red boxes. Data are the annual mean output filtered by

120 (left column) 10-year high pass, (middle column) 10-year to 50-year band pass and (right

121 column) 50-year low pass. (a), (b) and (c) are results from the Coupled, SlabOcean and

122 ClimSST simulations, respectively. Contour interval is 0.2. Dashed contours represent

123 negative values. The zero contour is thickened. Variance explained by the first EOF is

124 shown in the title of each panel. (d) shows the power spectra of the first PC (scaled by a

125 factor of 1/1000) for the corresponding timescales. It can be seen from (d.3) that the

126 multi-decadal “ENSO-like” variability in the SlabOcean run is stronger than that in the
127 Coupled run.

128

129 **Reference**

130 Hurrell, J. W., J. J. Hack, D. Shea, J. M. Caron, and J. Rosinski, 2008: A new sea surface
131 temperature and sea Ice boundary dataset for the community atmosphere model.
132 *J. Clim.*, **21**, 5145–5153, doi:10.1175/2008JCLI2292.1.

133